



FOGSI Position Statement

Vaccination in Pregnant Indian Women

CONTRIBUTORS

1. Dr. Ameya Purandare

MD, DNB, FCPS, DGO, DFP, MNAMS, FICMCH,
FICOG, Fellowship in Gyn Endoscopy
(Germany)

Consultant Obstetrician & Gynaecologist,
Purandare Hospital, K. J. Somaiya Medical
College and Super Specialty Hospital,
Sir H. N. Reliance Hospital, Bhatia Hospital,
Masina Hospital, Naigaon Municipal
Maternity Home, Mumbai Police Hospital,
Mumbai

Joint Treasure, FOGSI, 2026
President, AMOGS, 2026–28

2. Dr. Anahita Chauhan

MD, DGO, DFP, FICOG
Honorary Consultant,
Saifee, Bhatia and St Elizabeth Hospitals,
Mumbai
Joint Associate Editor, JOGI

3. Dr. Anju Soni

MBBS, MS
Senior Consultant and Head of the
Department, Obstetrics & Gynaecology,
Soni Group of Hospitals and Soni Manipal
Hospital, Jaipur
Vice President, West Zone, FOGSI, 2026

4. Dr. Asha Jain

MD, DNB
Consultant Obstetrician and Gynaecologist,
Director, SMSA Hospital, Raipur
Committee Chairperson, FDMSE Committee,
FOGSI

5. Dr. Bhaskar Pal

DGO, MD, DNB, FICOG, FRCOG
Senior Consultant, Obstetrics and
Gynaecology,
Apollo Multispeciality Hospital, Kolkata
President, FOGSI, 2026

6. Dr. Bidisha Roy Choudhury

Assistant Professor,
IPGMER, SSKM Hospital, Kolkata

7. Dr. Dibyendu Banerjee

DGO, MD, DNB, FICOG
Head of the Department, Department of
Gynecology and Obstetrics,
Charnock Hospital, Kolkata
Founder Partner, Ankuran Pluro Infertility and
IVF, Kolkata

8. Dr. Gayathri Dinesh Kamath

MD, MRCOG, FMAS
Additional Director, Department of Obstetrics
and Gynaecology,
Fortis Hospital, Bangalore

9. Dr. Girija Wagh

Head of Department of OBGYN, Bharati
Vidyapeeth Deemed to be University Medical
College, Pune
Director, Primrose Wellness and Care, Pune
Consultant, Cloudnine Hospitals and Apollo
Spectra Hospital, Pune
President Elect, Gestosis India Association
Member in the Steering Committee, World
Organisation Gestosis

10. Dr. Madhuri A. Patel

MD, DGO, FICOG
President Elect, FOGSI
Secretary General, FOGSI, June 2021–24

11. Dr. Meena Samant

MD, DNB, MRCOG
Director and Head of the Department,
Department of Obstetrics and Gynaecology,
Jay Prabha Medanta Hospital, Patna

12. Dr. Meenu Agarwal

Director and Consultant,
Morpheus Bliss Fertility Centre, Pune
Former President, Pune Obstetrics and
Gynaecological Society
Vice President, West Zone, FOGSI, 2026

13. Dr. Meeta Singh

Co-director, Head Ob-Gyn,
Tanvir Hospital, Hyderabad

14. Dr. Niranjan Chavan

Professor and Unit Chief, LTMMC & LTMGH,
Sion Hospital, Mumbai
Medical Director, Chavan Maternity &
Nursing Home, Mumbai
Treasurer, FOGSI, 2025–27
Past President, MOGS, 2022–23
President Elect, AFG, 2026–27
Vice Chairperson, Maharashtra Chapter of
IAGE, 2025–27
Organising Secretary, AICOG, Mumbai, 2025

15. Dr. Parikshit Tank

MD, DNB, FCPS, DGO, DFP, MNAMS, FICOG,
FRCOG
Co-founder, Pluro Fertility and IVF
Consultant Obstetrician and Gynecologist,
Ashwini Pluro Clinic and Center for
Endoscopy and IVF
Consultant, Zynova, Surya and Jupiter
Hospitals, Mumbai
Deputy Secretary General, FOGSI

16. Dr. Poushali Sanyal

MS, DNB, FMAS, MRCOG, FICOG
Director,
Welcare Medical Centre, Kolkata

17. Dr. Priyanka Pipara

DGO, DNB, FICOG, MRCOG, MNAMS, FICRS
Consultant, Obstetrics & Gynaecology
Apollo Multispeciality Hospital, Kolkata

18. Dr. Rashmi Bagga

MD, FICOG
Professor, Obstetrics & Gynaecology,
Post Graduate Institute of Medical Education &
Research, Chandigarh

19. Dr. Rohan Palshetkar

MS, FRM, BDRME, ADRME
Chairperson, FOGSI Young Talent Promotion
Committee
Professor, DY Patil School of Medicine
Head of Unit, Bloom IVF
Consultant, Sir H. N. Reliance Hospital, Saifee
Hospital, Surya Hospital, Palshetkar Patil
Nursing Home, Mumbai

20. Dr. Seetha Ramamurthy Pal

DGO, MD, FICOG, FRCOG
Senior Consultant, Obstetrics and Fetal
Medicine,
Apollo Multispeciality Hospital, Kolkata

21. Dr. Soma Datta

MS, FICOG, FMAS
Consultant Gynaecologist,
Woodlands Hospital, Kolkata
Joint Secretary, FOGSI, 2026

22. Dr. Suvarna Khadilkar

Consultant Obstetrician and Gynaecologist,
Mumbai
Secretary General, FOGSI

FOGSI Position Statement

Vaccination in Pregnant Indian Women



Background

Infections contribute substantially to maternal and neonatal morbidity and mortality; hence, maternal vaccination has significant public health relevance by reducing vaccine preventable diseases in the mother and the newborn.

Safety considerations for inactivated vs live vaccines

Available evidence indicates that, in general, inactivated vaccines can be safely administered during pregnancy, whereas live vaccines are avoided except in situations where the risk of disease exposure is substantial and benefits outweigh potential risks.¹

Pre-conceptual Vaccines

MMR or Rubella vaccine needs to be given to those who have not received the vaccine in adolescence. The MMR vaccine is preferred because of wider immunization coverage. Varicella vaccine should also be administered to those who have neither been infected nor vaccinated before.¹

Timing of Vaccination

- ▶▶ Pregnancy should be avoided for 1 month after vaccination.^{1,2}
- ▶▶ However, if the patient does conceive, she should be counselled regarding the theoretical risks to the fetus; however, it is not an indication for termination of pregnancy.^{1,2}

Vaccines Recommended During Pregnancy

Td, Tdap, and influenza vaccines are strongly recommended during each pregnancy.^{2,3}

Td vaccine (1st dose)

- ▶▶ Td (tetanus and diphtheria) has replaced TT since 2018.
- ▶▶ First dose should be given early in pregnancy (at or after 12 weeks).
- ▶▶ It is not required if the woman has been previously immunized with two doses of Td/TT during the previous pregnancy within the past 3 years.

Tdap vaccine

- ▶▶ Tdap vaccine is a combination vaccine that protects against three bacterial infections – tetanus, diphtheria and pertussis with a single dose.
- ▶▶ Neonatal morbidity and mortality due to pertussis is high in the first 2 months of life. Maternal Tdap vaccination covers the immunity gap till the neonate is vaccinated.²⁻⁴
- ▶▶ One dose between 27 and 36 weeks of pregnancy is recommended in each pregnancy regardless of prior vaccination history. (It should be given even if the woman has taken two doses of TT/Td in current pregnancy.)
- ▶▶ It should be considered in place of second dose of Td to provide protection against pertussis in addition to tetanus and diphtheria.^{3,4}
- ▶▶ If Tdap is not given during pregnancy, it should be given postpartum.⁴

Influenza vaccine

- ▶▶ Pregnant women are at increased risk of severe influenza infection, including pneumonia, hospitalization, ICU admission, and maternal mortality.^{4,5}
- ▶▶ Influenza infection during pregnancy has also been associated with adverse fetal outcomes, including early pregnancy loss, preterm birth, low birth weight, and stillbirth.
- ▶▶ One dose of inactivated influenza vaccine (quadrivalent or trivalent) is recommended at or after 12 weeks of pregnancy (earlier during pandemic)^{2,4,6} or 12 months after last dose (whichever is later).
- ▶▶ Administer the currently available strain (do not delay by awaiting newer strains).
- ▶▶ Maternal vaccination offers passive immunity to the newborn.
- ▶▶ Live attenuated influenza vaccine (intranasal) is contraindicated in pregnancy due to theoretical risk of fetal transmission.⁵

Vaccines Recommended during Pregnancy

FOGSI Position Statement

- ▶▶ First dose of Td should be given early in pregnancy (at or after 12 weeks).
- ▶▶ Tdap is the preferred tetanus-containing vaccine (in place of second dose of Td) during pregnancy and should be administered in the early third trimester, ideally between 27 and 36 weeks of gestation, to optimize passive antibody transfer and protect the newborn during the neonatal period.
- ▶▶ A single dose of Tdap is recommended during each pregnancy, irrespective of prior Tdap vaccination history.
- ▶▶ Women who are pregnant should receive one dose of inactivated influenza vaccine (IIV) or recombinant influenza vaccine (RIV).

Vaccines Contraindicated in Pregnancy

All live vaccines such as MMR, Varicella, HPV (Human Papilloma Virus), Intranasal influenza, BCG, and Japanese Encephalitis vaccines are contraindicated during pregnancy due to risk of fetal transmission.⁶

Vaccines Recommended in Special Situations

Vaccines may be considered in special situations when there is a risk of maternal infection and potential vertical transmission to the fetus and where the benefits outweigh the risks.

COVID-19 vaccine

- ▶▶ Pregnant women are at increased risk of adverse COVID-19 outcomes, including higher rates of ICU admission and adverse pregnancy outcomes.^{2,5}
- ▶▶ COVID-19 vaccination reduces the risk of severe disease in pregnant individuals and induces antibody response in the infant.
- ▶▶ Maternal vaccination has also been associated with reduced COVID-19-related hospitalization in infants younger than 6 months.³
- ▶▶ mRNA-based COVID-19 vaccines are recommended in any trimester of pregnancy.
- ▶▶ During a pandemic or local epidemic, COVID-19 vaccination is recommended for all eligible individuals, including pregnant women. A two-dose series should be administered, with doses given at least 4 weeks apart.²
- ▶▶ Booster doses may be considered during periods of increased viral transmission.

Hepatitis A vaccine

- ▶▶ Inactivated vaccine indicated in women at high risk of infection, including those with chronic liver disease, poor sanitation, exposure or travel to endemic areas.^{1,3}
- ▶▶ Vaccination provides protection within 2 weeks for up to 10-30 years.⁷

Hepatitis B vaccine

- ▶▶ Hepatitis B vaccination (inactivated or recombinant) is recommended during pregnancy in women at high risk of exposure.
- ▶▶ These include women with HBsAg-positive partners, multiple sexual partners, recent sexually transmitted infections, IV drug users, those receiving regular blood products, those with chronic liver or kidney disease and occupational exposure.

Rabies vaccine (live attenuated)

Post-exposure prophylaxis (PEP) with vaccination is recommended in pregnancy and lactation as benefits outweigh the risks.¹

Typhoid vaccine

Live vaccines are contraindicated in pregnancy; however, in clearly indicated situations, the inactivated Vi polysaccharide vaccine may be administered.^{1,8}

Yellow fever vaccine

- ▶▶ Vaccination may be considered if travel to endemic regions (e.g., Africa or South America) poses a significant risk of infection.^{1,8}
- ▶▶ Travel should ideally be postponed during pregnancy. If unavoidable, a single dose may be administered after careful evaluation of benefit vs risk or waiver may be sought.

Emerging Vaccines

Respiratory syncytial virus vaccine

- ▶▶ A single dose of respiratory syncytial virus (RSV) vaccine (inactivated) is advised in the third trimester of pregnancy.²
- ▶▶ It can be given to those who do not have a delivery planned in the next 2 weeks to prevent RSV-related lower respiratory tract infection (LRTI) in newborns.

Meningococcal vaccine

Vaccination decisions during pregnancy should be based on individual exposure risk and clinical judgment.²

Postpartum Vaccination

- ▶▶ The postpartum period is ideally suited for opportunistic vaccination.
- ▶▶ Vaccines which were not received in pre-conception or during pregnancy should be given in the postpartum period, if indicated. This include MMR, Varicella, Influenza, Hepatitis B, and Tdap vaccines.^{2,3}
- ▶▶ HPV vaccination schedule should be completed if interrupted by pregnancy or initiated postpartum.¹

Vaccines recommended during the pre-conception, pregnancy, and postpartum periods²

Pre-conception	Pregnancy	Postpartum
<ul style="list-style-type: none">▶▶ MMR▶▶ Varicella▶▶ Inactivated influenza vaccine, seasonally▶▶ Hepatitis B (if at risk and not vaccinated before)	<ul style="list-style-type: none">▶▶ Td (early in pregnancy, at or after 12 weeks)▶▶ Tdap (between 27 and 36 weeks)▶▶ Influenza (at or after 12 weeks, earlier during pandemic)	<ul style="list-style-type: none">▶▶ MMR (if at risk)▶▶ Varicella (if at risk)▶▶ Influenza▶▶ Hepatitis B▶▶ Tdap▶▶ HPV

References

1. World Health Organization (2025). *Immunization in pregnancy: WHO guidelines on vaccination for pregnant women*. Available at: <https://www.who.int/healthtopics/immunization>
2. Centers for Disease Control and Prevention (2025). *Vaccination during pregnancy*. Available at: <https://www.cdc.gov/vaccines/pregnancy/index.html>
3. Federation of Obstetric and Gynaecological Societies of India (FOGSI), International Clinical Obstetric Gynaecology (ICOG). *Vaccination in Women [Internet]*. 2024 Apr 26 [cited 2026 Feb 10]. Available at: https://www.fogsi.org/wpcontent/uploads/2024/08/Binder_Vaccination-in-Women.pdf
4. Mathur G, Vora A, et al.; for the Association of Physicians of India (API), RSSDI, ISN, GSI, IMA, FOGSI, ICS, IAPSM, & other collaborating societies. *Indian Consensus Guideline on Adult Immunization*. Mumbai: Association of Physicians of India; 2024. Available at: <https://apiindia.org/reader/Indian%20Consensus%20Guideline%20on%20Adult%20Immunization>
5. American College of Obstetricians and Gynecologists (ACOG) (2022). *ACOG Practice Bulletin No. 225: Immunization during Pregnancy*. *Obstetrics & Gynecology*, 139(3), pp. e1–e13. Available at: <https://doi.org/10.1097/AOG.0000000000004746>
6. Centers for Disease Control and Prevention (CDC) (2023). *Vaccines and Pregnancy*. Available at: <https://www.cdc.gov/vaccines/pregnancy.html>
7. Association of Obstetricians & Gynaecologists of Delhi. *AOGD Bulletin*. Vol. 23, Issue 11 [Internet]. New Delhi: Association of Obstetricians & Gynaecologists of Delhi; 2024 Apr [cited 2026 Feb 12]. Available at: <https://aogd.org/wpcontent/uploads/2025/04/AOGD-Bulletin-April-2024.pdf>
8. American College of Obstetricians and Gynecologists (ACOG) (2021). *Committee Opinion No. 736: Maternal Immunization*. *Obstetrics & Gynecology*, 137(3), pp. e45–e53. Available at: <https://doi.org/10.1097/AOG.0000000000004116>